

Biochemistry of Skin

Assoc.Prof.Siriwan Ongchai, Ph.D

Department of Biochemistry,
Faculty of Medicine,
Chiang Mai University.

Objective

☀ **Structure and Function of skin biomolecules**


☀ **Skin junction and diseases**

☀ **Skin Melanogenesis: mechanism and functions**


☀ **Skin Melanogenesis and associated disorders**

☀ **skin ageing**

☀ **Nutrition and skin**



Functions of Skin



Functions of Skin

Protection:

Sensation:



Heat regulation:

Control of evaporation:

Storage and synthesis:

Excretion: Absorption:

Water resistance:



This figure shows the conversion of 7-dehydrocholesterol to vitamin D in the skin and its transfer to the blood where it is complexed to a Vitamin D binding protein (DBP).

Epidermal Biomeolecules


Intercellular spaces between keratinocytes

**Lipid leaflets
(barrier lipid)** → **Glycolipids, Sterols, Phospholipids**

Proteoglycans →

Epican; HS/CS proteoglycan,


Epican is a heparan/chondroitin sulfate proteoglycan form of CD44 and is expressed on the surface of keratinocytes from the basal layer to the granular layer of the epidermis.




One way to understand the barrier function of the stratum corneum is to consider it as a brick wall. The corneocytes (made of tough protein) form the bricks and between these a double layer of lipids (fatty materials) and water make up the mortar. Some lipids have a hard crystal-like structure and are impermeable to water. Others lipids do not have this structure and they allow water to percolate through. So, the barrier is semi-permeable.

<http://www.hse.gov.uk/skin/professional/causes/understand.htm>

Dermal Biomeolecules (Dermal ECM)




Dermal Biomeolecules (Dermal ECM)

Fibrous protein:  Collagens
Elastin

Ground substance:

Glycosaminoglycans → Hyaluronan

Proteoglycan  Small DSPG
Large CSPG

Glycoproteins and enzymes; MMPs




Dermal collagens: a major component of skin

type I ~ 85-90%

type III ~ 8-11% (more prominent in fetal skin and early stage of wound healing)

type V ~ 2-4%

small proportions; type VI, XII, XIII



Copyright © 2000 Benjamin/Cummings, an imprint of Addison Wesley Longman, Inc.



Collagen peptide

It is also called collagen hydrolysate, gelatine, gelatine hydrolysate and hydrolyzed gelatine.

Collagen peptide biological activities

Bovine hide CH(GELTA[®] CPB1000) stimulated cell proliferation, *HAS2* mRNA expression and hyaluronan production in human dermal fibroblasts

J Dermatol. 37(4): p. 330-8.

After 4 weeks of daily ingestion of CH, it promoted stimulatory effects on skin tissue, increasing expression levels of type I and type IV collagen

J Med Food. 14(6): p. 618-24.

Collagen hydrolysate ingestion inhibited ultraviolet B-induced decreased of type I collagen, thus improving skin conditions in mice

13

Biosci Biotechnol Biochem, 2009. 73(4): p. 930-2.


Side Effects of Taking Collagen Supplements

- **High Calcium Levels** → hypercalcemia.

Too much calcium in the body causes constipation, bone pain, fatigue, nausea, vomiting and abnormal heart rhythms.


- **Hypersensitivity Reactions** → abnormal response of the immune system against allergens

Dermal Biomeolecules (Dermal ECM)

Fibrous protein:  Collagens
Elastin


Ground substance:

Glycosaminoglycans → Hyaluronan


Proteoglycan  Small DSPG
Large CSPG

Glycoproteins and enzymes; MMPs






Dermal Biomeolecules (Dermal ECM)

Fibrous protein:  Collagens
Elastin

Ground substance:

Glycosaminoglycans → Hyaluronan

Proteoglycan  Small DSPG
Large CSPG


Glycoproteins and enzymes; MMPs



Glycosaminoglycans (GAGs)

***Jelly-like property**

*** Heteropolysaccharide**




***No sulfate in the chain**


***No covalent bonding with protein to form proteoglycan**

*** High content in child and gradually decrease with age**

Loss of HA associated with skin aging




Skin with high HA content



Skin with low HA content




Bacterial Hyaluronidase

Bacterial hyaluronidases, enzymes capable of breaking down hyaluronate, are produced by a number of pathogenic Gram-positive bacteria that initiate infections at the skin or mucosal surfaces.




Proteoglycan)


Decorin



Decorin knock-out mice



Hypodermis



The hypodermis is the innermost and thickest layer of the skin. It invaginates into the dermis and is attached to the latter, immediately above it, by collagen and elastin fibres.

It is essentially composed of a type of cells specialized in accumulating and storing fats, known as adipocytes.

These cells are grouped together in lobules separated by connective tissue .

Objective

☀ **Structure and Function of skin biomolecules**


☀ **Skin junction and diseases**

☀ **Skin Melanogenesis: mechanism and functions**

☀ **Skin Melanogenesis and associated disorders**

☀ **skin ageing**


☀ **Nutrition and skin**




Skin Junction

- **Dermoepidermal Junction**
 - ☐ **Hemidesmosomal Junction**
 - ☐ **Desmosomal Junction**



Skin Junction




Dermoepidermal Junction



Basement Membrane





Basal membrane (Basement Membrane)


Epidermolysis bullosa dystrophica (Dystrophic EB; DEB)


- **The deficiency in anchoring fibrils impairs the adherence between the epidermis and the underlying dermis.**
- **The skin of DEB patients is thus highly susceptible to severe blistering.**
- **Open wounds on the skin heal slowly or not at all, and are particularly susceptible to infection.**
- **The chronic inflammation leads to errors in the DNA of the affected skin cells, which in turn causes squamous cell carcinoma (SCC). The majority of these patients die before the age of 30.**

Epidermolysis bullosa dystrophica (**Dystrophic EB**; DEB)



Skin Junction







Bullous pemphigoid

- Bullous pemphigoid is characterized by the presence of immunoglobulin autoantibodies specific for the hemidesmosomal bullous pemphigoid antigens BP230 (BPAg1) and BP180 (BPAg2).

Skin Junction



Desmosomal Junction




Disturbed desmosomal adhesion contributes to the pathogenesis of a number of diseases such as pemphigus vulgaris, which is caused by autoantibodies against desmosomal cadherins or viral infection such as herpes simplex infections.

Cosmetic applications of skin biomolecules: anti-wrinkle



Most fillers replace the host components of the skin: collagen, hyaluronic acid, and elastin



Collagen: collagen makes up 75% of our skin protein. From around the age of 25 Collagen levels start to deplete at an astonishing rate of 1.5% every year.

Bovine Collagen Fillers

Bovine collagen is processed from the skin of cows. Approved in the 1980s as a wrinkle treatment, bovine collagen is still widely used as a cosmetic filler.

It can cause allergic reactions, The body naturally breaks down injected collagen, so you need to get collagen injections two to four times per year to maintain results.

Human Collagen Fillers

Human collagen, made from cultures of human cells, became available in 2002. Human collagen causes dramatically fewer allergic reactions than bovine collagen. It is more expensive than bovine collagen, and injections also need to be repeated every three to six months.


Hyaluronic acid:

its molecules bind to water in the skin, hydrating and firming its structure, indicate that HA injections induce collagen production, stimulation of growth factors and inhibition of collagen breakdown, lasting for 4-12 month

Allergic reactions are very rare.

Fat Injection Fillers

Fat injections involve removing small amounts of fat from the thighs, belly, or buttocks and injecting it under the skin of the face. The fat expands the skin, shrinking wrinkles. Because it is your own tissue, there can be no allergic reaction.





Derived from Botulinum toxin type A, **Botox** blocks nerve impulses within facial muscles, thus smoothing wrinkles by temporarily paralyzing the muscles. It is





ing use
brows
to alw
nce of
It also
the e
sfully c
all of th
ten mi
last a



deep lines
ead lines that
ed. It reduces
r at the side
scler which
me doctors
to lessen the
The process
ections, with
ths.

etic is a protein that is injected into
relax them, smoothing lines and wrinkles
peated facial movements.

Skin Pigmentation (Melanogenesis)




Skin Pigmentation (Melanogenesis)


MC1R
(melanocortin-1
receptor)


MITF
(microphthalmia
transcription factor)


TRYP1:Tyrosinase-
related protein 1



Skin Pigmentation (Melanogenesis)








Simplified Pathway of Melanin Formation

Illustrated by USHP








FIGURE 1. Schematic of human skin architecture from light- and dark-pigmented skin types. From top to bottom: SC, stratum corneum; G, stratum granulosum; S, stratum spinosum; B, stratum basale; BM, basement membrane; D, dermis. Cell types: K, keratinocyte; M, melanocyte; F, fibroblast; shaded oval, melanin granule


Complexion and biology

	AFRICAN	ORIENTAL	CAUCASIAN
Type of Melanin Mixture	Pheomelanine Eumelanine	Pheomelanine Eumelanine	Pheomelanine Eumelanine
Proportion of Melanin Grains in the Epidermis	Complexed Free	Complexed Free	Complexed Free
Melanine Morphology Grain			
Melanine Grain Size (nm)	1 x 0.5	0.6 x 0.3	0.5 x 0.3




Target mechanism of Whitening agents






endocytotic activity of keratinocyte




niacinamide

a Immediate pigment darkening (IPD)




b Delayed tanning (DT)





The mechanism of skin pigmentation after UV irradiation


Expert Reviews in Molecular Medicine©2002 Cambridge University Press



DCT=dopachrome tautomerase
Inhibition by Licorice, Kogic acid, Vit C

TRYP1:Tyrosinase-related protein 1





Biological Function of Melanin

Melanins function to absorb and disperse UV light and act as free radical scavengers, thus reducing the penetration of UV radiation.

Melanogenesis also serves as a major antioxidant defence mechanism of skin as melanocytes, the specialised cells that produce melanins, neutralise the deleterious effects of active oxygen species and free radicals.

Hyperpigmentation

- Increase in MC-1R on melanocyte membrane
- Increase in MC-1R and melanocyte proliferation
- Activation the mechanism of action of melanogenesis

Hyperpigmentation related diseases

- Acanthosis nigricans
- Cushing's disease
- Addison's disease
- Linea nigra
- Pellagra
- Some cancer drug: cisplatin



Hyperpigmentation in
Acanthosis nigricans
(insulin resistance)



Hyperpigmentation in
Addison's disease
(increase synthesis MSH)



Hyperpigmentation in
Cushing's syndrome
(increase synthesis MSH)





Linea nigra

Post Inflammatory Hyperpigmentation (PIH)




**Post Inflammatory Hyperpigmentation (PIH)
after Photodynamic Therapy treatment for acne.**


Hypopigmentation

- Decrease in the number of melanocyte
- Lack of tyrosine

Metabolic Pathways of Phenylalanine




*Phenylalanine hydroxylase (PAH)



* tyrosinase

Metabolic Pathways of Phenylalanine




Biochemistry of Skin Aging




✧ Intrinsic Aging

✧ Extrinsic or photo aging



20s

Free Radicals Attack Surface
Environmental Damage is High




Collagen
Fibres Healthy Vascular
Tissue

Constant exposure to the sun
& fast life style leads to skins
premature aging.

30s

First Signs of Aging Appear
Dull, Lackless, Lifeless Skin




Collagen
Fibres Lessening Vascular
Tissue

Skin regeneration is reducing
leading to dull complexion &
uneven skin tone. Use of harsh
skin care can become apparent.
Elastin degradation can show
first signs of aging.

40s

Significant dullness, aging &
dark spots. Skin sensitivity.




Collagen
Fibres Reducing Vascular
Tissue

Skin thinning can cause
sensitivity, redness, dry, oily,
sudden redness. Photo aging
appears - dark spots - more
prominent signs of aging
appear.


50+

Significant tension decrease.
Moisture retention slowing.



Collagen
Fibres Reduced Vascular
Tissue Thin

Decrease in surface tension impairs
skin structure and ability to defend
itself. Barrier lessens leading to
less efficiency in retaining moisture.
Combined with excessive dryness
sometimes accompanied by adult acne.



Biochemistry of Skin Aging

Effects on cells → Decrease in cell apoptosis





Effects on skin ECM


Effects on proteins

Effects on lipids


Effects on DNA

Effects on proteoglycans and *GAGs*





Advanced Glycation Ends products
(AGEs)






Nutrition and Skin



EFA



Water



Minerals



Proteins



Vitamins





Pellagra:
Niacin def.



Vitamin B3, also known as Niacin or Nicotinic acid, is a water-soluble B-vitamin that plays a significant role in energy metabolism and maintaining healthy skin .

Niacin is synthesized inside the body by the liver from the essential amino acid tryptophan.


Niacin is lost with cooking.
Also, alcohol inhibits vitamin B3 absorption.

*Corn is the only grain low in niacin.



Fig. 1: Principle Action of B3 Metabolism in the Body

Tryptophan



Deficiency of vitamin B (Vit B2, Vit B5, Vit B9 or vit B12)

- hair changes, and recurrent angular stomatitis





Hair loss (alopecia)
caused by biotin (vit B7) deficiency




Fig. 6-6 Alopecia before (A) and after (B) biotin therapy in a patient

Objective

☀ **Structure and Function of skin biomolecules**

☀ **Skin junction and diseases**

☀ **Skin Melanogenesis: mechanism and functions**

☀ **Skin Melanogenesis and associated disorders**

☀ **skin ageing**

☀ **Nutrition and skin**

